

Note to Readers: If you need assistance accessing items in this Supplemental Material, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Table of Contents for Supplemental Material

Biased Exposure–Health Effect Estimates from Selection in Cohort Studies: Are Environmental Studies at Particular Risk?

Marc G. Weisskopf, David Sparrow, Howard Hu, and Melinda C. Power

Methods

Details of inverse probability weighting

Table S1. Variables used in forward selection logistic regression model to calculate inverse probability of attrition weights (IPW).

Table S2. Variables included in the final inverse probability of attrition weighting model.

Table S3. Characteristics at the time of bone lead measurement among all those with bone lead measurement (N=835).

Table S4. Fully adjusted^a hazard ratios (95% confidence intervals) for all-cause, cardiovascular disease, and ischemic heart disease mortality, by tertile^b of patella lead at baseline among white men in the Normative Aging 45 years old or younger at NAS study entry (N=637), and applying inverse probability of attrition (IPW) weights truncated at the 1st and 99th percentile of the distribution of the IPW weights.

Table S5. Adjusted hazard ratios (HR; 95% CI) for all-cause, cardiovascular disease, and ischemic heart disease mortality, by tertile^b of blood lead at baseline among either all white men in the Normative Aging Study (N=1,206), or those 45 years old or younger at NAS study entry (N=909).

Table S6. Adjusted hazard ratios (HR; 95% CI) for all-cause, cardiovascular disease, and ischemic heart disease mortality, by tertile of tibia lead at baseline among either all white men in the Normative Aging Study (N=834), or those 45 years old or younger at NAS study entry (N=636).

Figure S1. Nonlinear association between patella bone lead concentration and the log of HR (logHR) for all-cause, cardiovascular, and ischemic heart disease adjusted for age at KXRF, age at KXRF squared, smoking (never/former/current & packyears), and education among all white men (n=835) (Model 1: Base Model). The reference logHR=0 is at the mean of patella lead concentration. The estimates are indicated by the solid line and the 95% CIs by the dashed lines. The *P* values for significance of the nonlinear component for all-cause, cardiovascular, and ischemic heart disease mortality were 0.39, 0.54, and 0.64, respectively. Patella lead concentrations of all individual participants are indicated by short vertical lines on the x-axis..

Figure S2. Nonlinear association between patella bone lead concentration and the log of HR (logHR) for all-cause, cardiovascular, and ischemic heart disease adjusted for age at KXRF, age at KXRF squared, smoking (never/former/current & packyears), and education among white men 45 years old or younger at NAS entry (n=637) and with inverse probability weighting to weight the analyses to reflect the full group still alive at the time of KXRF (Model 4). The reference logHR=0 is at the mean of patella lead concentration. The estimates are indicated by the solid line and the 95% CIs by the dashed lines. The *P* values for significance of the nonlinear component for all-cause, cardiovascular, and ischemic heart disease mortality were 0.48, 0.91, and 0.28, respectively. Patella lead concentrations of all individual participants are indicated by short vertical lines on the x-axis.